

DUFACOX 2.5% ORAL

Composition:

Contains per ml solution:

Toltrazuril..... 25 mg

Description:

Dufacox 2.5% Oral is a anticoccidial which is active against all intracellular development stages of coccidia. Toltrazuril is effective in the treatment of coccidiosis in poultry, turkeys, calves, lambs, kids and piglets caused by different species of *Eimeria spp.*

Indications:

Coccidiosis of all stages like schizogony and gametogony stages of different *Eimeria spp.* in chickens, turkeys, calves, lambs, kids and piglets.

Contraindications:

Do not administer to animal with an impaired liver and/or renal function.

Side effects:

At high dosages in laying hens egg-drop and in broilers growth inhibition and polyneuritis can occur.

Dosage and administration:

For oral administration via drinking water.

Chickens and turkeys: 1 litre per 1000 litres of drinking water continuous medication for 2 days (7.5 mg/kg body weight)

Calves, lambs, kids and piglets: 0.8 ml per kg body weight for 2 days (20 mg/kg body weight).

During treatment the medicated water should be the only source of drinking water!

Medicated water should be used immediately and refreshed every 24 hours for the duration of the treatment. The body weight should be determined with accuracy.

Withdrawal times:

Meat and offal: 19 days (poultry and turkeys)

42 days (sheep and goats)

63 days (calves)

77 days (pigs)

Do not use in breeders in production or layers producing eggs for human consumption.

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

100 ml bottles and 1 litre cans.