

## **DUFAXACIN 10% ORAL**

### **Composition:**

Contains per ml oral solution:

Enrofloxacin ..... 100 mg

### **Description:**

Enrofloxacin is a fluoroquinolone antibiotic with bactericidal activity against both gram-positive and gram-negative bacteria as well as *Mycoplasma* spp, *Chlamydia* spp. and *Rickettsia* spp.

### **Indications:**

Gastrointestinal and respiratory infections in swine and poultry, caused by microorganisms susceptible to enrofloxacin, such as *Campylobacter* spp., *E. coli*, *Haemophilus* spp., *Mycoplasma* spp., *Pasteurella* spp. and *Salmonella* spp.

### **Contraindications:**

Hypersensitivity to quinolones, severe hepatic or renal impairment.

### **Dosage and administration:**

For oral administration in drinking water.

Swine ..... 1 ml per 40 kg body weight

(2.5 mg enrofloxacin per kg body weight)

Poultry ..... 1 litre per 2,000 litres of water for 3-5 days

(10 mg enrofloxacin per kg body weight)

The body weight should be determined with accuracy. Shake well before use. Medicated water should be used immediately and refreshed every 24 hours for the duration of the treatment.

### **Side effects:**

Gastrointestinal disturbances.

### **Precautions and warnings:**

It is recommended that the susceptibility of the causative microorganisms is established before treatment is commenced.

Severely sick animals have altered drinking patterns and may need to be medicated parenterally.

Dosage may need to be adjusted to reflect herd or flock water consumption.

### **Withdrawal times:**

Meat and offal: 4 days

Do not use in layers producing eggs for human consumption.

### **Storage conditions:**

Store dark and below 25°C. Do not freeze.

### **Packing:**

100 ml and 1 litre bottles.