

## **DUFAXACIN 10% INJ**

### **Composition:**

Contains per ml solution:

Enrofloxacin ..... 100 mg

### **Description:**

Enrofloxacin is a fluoroquinolone antibiotic with bactericidal activity against both gram-positive and gram-negative bacteria as well as *Mycoplasma* spp, *Chlamydia* spp. and *Rickettsia* spp.

### **Indications:**

Gastrointestinal and respiratory infections such as pasteurellosis, mycoplasmosis, colibacillosis, colisepticemia and salmonellosis in cattle and swine. Secondary bacterial infections during the course of viral disease, such as bovine respiratory disease complex.

### **Contraindications:**

Hypersensitivity to quinolones, severe renal or hepatic impairment.

### **Dosage and administration:**

For subcutaneous (cattle) or intramuscular (swine) administration.

Recommended dose: 1 ml per 40 kg body weight per day for 3 days

In severe cases of respiratory infections and salmonellosis: up to 2 ml per 40 kg body weight per day for 3 days (swine) or 5 days (cattle)

To avoid underdosing, the body weight of the animals should be determined with accuracy.

### **Side effects:**

Injection site reactions.

### **Precautions and warnings:**

It is recommended that the susceptibility of the causative microorganisms is established before treatment is commenced.

### **Withdrawal times:**

Meat and offal (cattle and swine): 10 days

Milk (cattle): 4 days

### **Storage conditions:**

Store dark and below 25°C. Do not freeze.

### **Packing:**

100 ml vials.