

DUFAMOX 15% LA INJ

Composition:

Contains per ml oily suspension:

Amoxicillin (as trihydrate) 150 mg

Description:

Amoxicillin is broad spectrum semi-synthetic penicillin with bactericidal activity against a wide range of gram-positive and gram-negative bacteria including the following:

Gram-negative: *Actinobacillus lignieresii*; *Bordetella bronchiseptica*, *Escherichia coli* (non β -lactamase producing strains), *Corynebacterium* spp.; *Fusobacterium* spp.; *Haemophilus* spp.; *Moraxella* spp.; *Pasteurella* spp.; *Proteus mirabilis* (non β -lactamase producing strains); *Salmonella* spp.

Gram-positive: *Bacillus anthracis*; *Clostridium* spp.; *Erysipelothrix rhusiopathiae*; streptococci; staphylococci (non β -lactamase producing strains).

Indications:

Bacterial infections in cattle, sheep, goats, pigs, dogs and cats caused by microorganisms susceptible to amoxicillin.

Contraindications:

Hypersensitivity to penicillins.

Dosage and administration:

For intramuscular injection in cattle, sheep, goats and pigs.

For subcutaneous injection in dogs and cats.

Recommended dose: 1 ml per 10 kg

If required, repeat after 2 days.

Shake well before use and always administer aseptically.

The dose volume should not exceed 20 ml in cattle, 10 ml in swine, 5 ml in sheep and goats at any one injection site.

To avoid underdosing, the body weight of the animals should be determined with accuracy.

Side effects:

Use of the product may occasionally result in local tissue reaction.

Precautions and warnings:

Therapy should be based on susceptibility testing or local epidemiological information. Do not handle the product if you are known to be hypersensitive to penicillins.

Withdrawal times:

Meat and offal: 14 days

Milk: 3 days

Storage conditions:

Store dark and below 25°C. Do not freeze.

After first opening the product should be stored dark and refrigerated (ca. 8°C).

Packing:

100 ml vials.