

DUFANOR 20% ORAL

Composition:

Contains per ml oral solution:

Norfloxacin..... 200 mg

Description:

Norfloxacin is a fluoroquinolone antibiotic with bactericidal activity mainly against gram-negative bacteria, as well as *Mycoplasma* spp.

Indications:

Gastrointestinal and respiratory infections in poultry caused by microorganisms susceptible to norfloxacin, such as Chronic Respiratory Disease (CRD), colibacillosis, Fowl Cholera, staphylococcal and streptococcal infections.

Contraindications:

Hypersensitivity to quinolones, severe hepatic or renal impairment.

Dosage and administration:

For oral administration in drinking water.

Recommended dose: 12 – 15 mg/kg body weight, equivalent to 250 ml per 1000 litres of water, continuously for 3-5 consecutive days

Shake well before use!

The consumption of drinking water should be determined with accuracy. Medicated water should be used immediately and refreshed for every 24 hours for the duration of treatment.

Side effects:

Gastrointestinal disturbances, excitation.

Precautions and warnings:

It is recommended that the susceptibility of the causative microorganisms is established before treatment is commenced.

Severely sick animals have altered drinking patterns and may need to be medicated parenterally.

Dosage may need to be adjusted to reflect herd or flock water consumption.

Withdrawal times:

Meat, offal and eggs: 4 days.

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

100 ml and 1 litre bottles.