

## **DUFAQUONE 5% INJ**

### **Composition:**

Contains per ml solution:

Buparvaquone ..... 50 mg

### **Description:**

Buparvaquone is a second generation hydroxynaphtaquinone active against both the schizont and piroplasm stages of *Theileria* spp., and can be used during the incubation period or when clinical signs are apparent.

### **Indications:**

Theileriosis in cattle, caused by the intracellular protozoan parasites *Theileria parva* (East Coast fever, Corridor disease, Zimbabwean theileriosis) and *T. annulata* (tropical theileriosis).

### **Contraindications:**

Hypersensitivity to buparvaquone. Do not administer by IV or SC injection. Theileriosis can interfere with the immune system, therefore a vaccination program should be postponed.

### **Dosage and administration:**

For intramuscular administration, preferable in the neck musculature.

Recommended dose..... 1 ml per 20 kg body weight

In severe cases the treatment may be repeated after 48-72 hours and the new injection should be administered on the other side of the neck, opposite to the first site of injection. The body weight should be determined with accuracy. Do not administer more than 10 ml in one injection site.

### **Side effects:**

Localised, painless, oedematous swelling may occasionally be seen at the injection site.

### **Withdrawal times:**

Meat and offal: 42 days

Milk: 2 days

### **Storage conditions:**

Store dark and below 25°C. Do not freeze.

### **Packing:**

50 ml vial.