

FENOSVIT-C W.S.P.

Composition:

Contains per g water soluble powder:

Colistin sulphate	200,000 IU	Thiamine HCl (B1)	2 mg
Erythromycin thiocyanate	35 mg	Riboflavin (B2)	4 mg
Neomycin sulphate	10 mg	Pyridoxine HCl (B6)	2 mg
Oxytetracycline HCl	50 mg	Cyanocobalamin (B12).....	10 mcg
Streptomycin sulphate	35 mg	Nicotinamide (PP, B3).....	20 mg
Vitamin A acetate.....	500 IU	Ascorbic acid (C).....	20 mg
Cholecalciferol (D3)	200 IU	Inositol	1 mg
Tocopherol acetate (E).....	2 mg	Calcium D-Pantothenate (B5)	10 mg
Menadione sodium bisulfite (K3)....	2 mg		

Indications:

Treatment and prevention of bacterial infections caused by gram-positive and gram-negative microorganisms and vitamin deficiencies due to infections and other stressors in poultry, swine and calves.

Contraindications:

Hypersensitivity to polymixins, macrolides, aminoglycosides or tetracyclines, severe hepatic or renal impairment.

Dosage and administration:

For oral administration in drinking water.

The recommended dosage levels are as followings:

- Therapeutic dosage: 1 g per litre of drinking water, continuously, for 5-7 consecutive days. In poultry this is equivalent to intake of 2.24 mg/kg b.w. colistin sulfate, 9.45 mg/kg b.w. erythromycin thiocyanate, 2.70 mg/kg b.w. neomycin sulfate, 13.5 mg/kg b.w. oxytetracycline hydrochloride and 9.45 mg/kg b.w. streptomycin sulfate.

- Prophylactic dosage: 0.5 g per litre of drinking water, continuously, for 5-7 consecutive days. In poultry this is equivalent to intake of 1.08 mg/kg b.w. colistin sulfate, 4.55 mg/kg b.w. erythromycin thiocyanate, 1.30 mg/kg b.w. neomycin sulfate, 6.50 mg/kg b.w. oxytetracycline hydrochloride and 4.55 mg/kg b.w. streptomycin sulfate.

The consumption of drinking water should be determined with accuracy. Medicated water should be used immediately and refreshed for every 24 hours for the duration of treatment.

Side effects:

Gastrointestinal disturbances.

Precautions and warnings:

It is recommended that the susceptibility of the causative microorganisms is established before treatment is commenced.

Severely sick animals have altered drinking patterns and may need to be medicated parenterally. Dosage may need to be adjusted to reflect herd or flock water consumption.

Withdrawal times:

Meat and offal: 10 days.

Eggs: 15 days.

Do not use in animals producing milk for human consumption.

Storage conditions:

Store dry and below 25°C. Do not freeze.

Packing:

100 g sachets and 1 kg jars.