

LIVER TONIC ORAL

Composition:

Contains per ml solution:

Acetyl methionine.....	17.4 mg	Riboflavin sodium phosphate ...	4.75 mcg
Choline chloride	10 mg	Thiamine HCl	0.95 mg
Cyanocobalamin	0.006 mcg	Biotin.....	2 mcg
Calcium pantothenate	581 mcg	Folic acid.....	10 mcg
Nicotinamide	1.1 mg	Ascorbic acid	35 mcg
Pyridoxine HCl.....	107 mcg	α-Tocopherol acetate	35 mcg

Description:

The liver functions as a processing plant, transforming, storing and detoxifying raw materials for the rest of the body. Liver impairment may lead to bleeding disorders, jaundice, reduction of growth and feed uptake, altered mental status and a wide range of secondary health issues such as skin disorders and coat abnormalities. Liver Tonic Oral is formulated to optimise liver function and aid in the prevention of toxic liver syndromes.

Indications:

An aid in the treatment and prevention of liver dysfunction and toxic liver syndromes in all animal species.

Contraindications:

None.

Dosage and administration:

For oral administration.

Large animals (≥100 kg) 5 ml per 100 kg body weight per day

Small animals (<100 kg) 1 ml per 10 kg body weight per day

In drinking water (incl. poultry)..... 1 litre per 2000 litres of drinking water, continuously

Once diluted, Liver Tonic Oral should be used immediately and refreshed every 24 hours for the duration of the treatment.

Withdrawal times:

None

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

1 litre bottles.