

OXYTETRACYCLINE 20% L.A. INJ

Composition:

Contains per ml solution:

Oxytetracycline (as dihydrate) 200 mg

Description:

Oxytetracycline is a highly active, broad spectrum tetracycline antibiotic with bacteriostatic activity against a wide variety of gram-positive and gram-negative organisms. The product is specifically formulated to provide a prolonged action resulting in sustained antibacterial activity.

Indications:

Bacterial infections in cattle, swine and sheep, caused by microorganisms susceptible to oxytetracycline such as atrophic rhinitis, navel/joint ill, respiratory tract infections, metritis, mastitis, urinary tract infections, shipping fever, liver abscesses, salmonellosis, dysentery, erysipelas, foot rot and the control of enzootic abortion in sheep.

Contraindications:

Hypersensitivity to tetracyclines, severe renal or hepatic impairment. Not for use in horses, dogs and cats.

Dosage and administration:

For intramuscular or subcutaneous administration.

Recommended dose: 1 ml per 10 kg body weight

If required, repeat after 48 hours.

The dose volume at one injection site should not exceed 20 ml in cattle, 10 ml in pigs or 5 ml in sheep.

To avoid underdosing, the body weight of the animals should be determined with accuracy.

Side effects:

Transient local injection site reactions. The use of oxytetracycline during the period of tooth and bone development, including late pregnancy may lead to tooth discoloration.

Withdrawal times:

Meat and offal: 21 days

Milk: 10 days

Storage conditions:

Store dark and below 25°C. Do not freeze.

Packing:

100 ml vials.