

BROILER FINISHER

Corn-soybased diet

Analytical constituents:

80.1%	Crude ash	16.9 %.....	Calcium	58.96 g/kg	Lysine
0.2 %	Crude fibre	0.83 %.....	Magnesium	86.10 g/kg	Methionine
13.5 %	Crude protein	4.89 %.....	Sodium	13.88 g/kg	Threonine
0.05 %	Crude fat	1.8 %.....	Phosphorus	0.01 g/kg	Tryptophan

Composition:

Calciumcarbonate, Monocalcium phosphate, Sodiumb carbonate, Sodiumchloride.

Additives (per kg)

Nutritional additives

Trace elements: 1,920 mg Fe (3b103 Iron(II)sulphate monohydrate); 64 mg I (3b202 Calcium iodate); 480 mg Cu (E4 Cupric(II)sulphate-pentahydrate); 2,240 mg Mn (3b502 Manganese(II)oxide); 2,240 mg Zn (3b605 Zinc sulphate monohydrate); 8 mg Se (E8 Sodium Selenite)

Vitamins: 384,000 IU Vitamin A (3a672a Retinyl acetate); 96,000 IU Vitamin D3 (3a671); 1,280 mg Vitamin E (3a700 dl-tocopherol); 80 mg Vitamin K3 (3a710 Menadione sodium bisulphite); 64 mg Vitamin B1 (3a821 Thiamine mononitrate); 192 mg Vitamin B2 (Riboflavin); 1,600 mg Vitamin B3 (3a315 Niacinamide); 480 mg Vitamin B5 (3a841 Calcium-d-pantothenate); 160 mg Vitamin B6 (3a831 Pyridoxine hydrochloride); 960 mcg Vitamin B12 (Cynacobalamin); 48 mg Vitamin B9 (3a316 Folic acid); 9,600 mg Vitamin B4 (3a890 Choline chloride); 4,800 mcg Vitamin H (3a880 Biotin)

Technological additives

Antioxidants: 160 mg Citric acid (E330); 53.3 mg B.H.T (E321); 53.3 mg Propyl gallate (E310).

Anti-caking agents: 150 g Clinoptilolite of sedimentary origin (1m558i); 40 mg Colloidal silica (E551b)

Zootechnical additives

Digestibility enhancers: 40,000 FYT 6-Phytase (4a18); 22,400 TXU Endo-1,4- β -xylanase (4a1602i); 10,000 TGU Endo-1,3(4)- β -glucanase (4a1602i)

Description:

Broilers have different feed requirements in terms of energy, proteins and minerals during different stages of their growth. It is important to adapt feed rations to these requirements for maximum production. As the broilers grow their energy requirements for the deposit of fat increase and their protein requirements decrease. Changes in body composition can be rapid during the finishing period and excessive fat deposition and loss of breast meat yield need to be carefully considered. DufaMix Broiler Finisher has been formulated to complement corn and soybean meal based diets for optimal performance and to allow the bird to express its full meat yield potential. Enzymes are included for improved feed conversion, digestion and the reduction of wet droppings.

Inclusion rate:

To be included in feed for broilers of approximately 28 days and older. The recommended inclusion rate for DufaMix Broiler Finisher is 25 kg per tonne of feed.

Storage conditions:

Store dry, dark and below 25°C. Do not freeze.

Packing:

10 and 25 kg sacks.