

## **BROILER STARTER**

## **Corn-soybased diet**

### **Analytical constituents:**

75.8 % .....	Crude ash	14.9 %.....	Calcium	67.49 g/kg .....	Lysine
0.51 % .....	Crude fibre	0.56 %.....	Magnesium	102.7 g/kg .....	Methionine
15.6 % .....	Crude protein	5.16 %.....	Sodium	18.17 g/kg .....	Threonine
0.34 % .....	Crude fat	6.82 %.....	Phosphorus	0.01 g/kg .....	Tryptophan

### **Composition:**

Monocalcium phosphate, Calciumcarbonate, Sodiumb carbonate, Sodiumchloride.

### Additives (per kg)

#### Nutritional additives

Trace elements: 2,600 mg Fe (3b103 Iron(II)sulphate monohydrate); 100 mg I (3b202 Calcium iodate); 600 mg Cu (E4 Cupric(II)sulphate-pentahydrate); 3,200 mg Mn (3b502 Manganese(II)oxide 3,200 mg Zn (3b605 Zinc sulphate monohydrate); 12 mg Se (E8 Sodium Selenite)  
Vitamins: 540,000 IU Vitamin A (3a672a Retinyl acetate); 160,000 IU Vitamin D3 (3a671); 2,800 mg Vitamin E (3a700 dl-tocopherol); 120 mg Vitamin K3 (3a710 Menadione sodium bisulphite); 140 mg Vitamin B1 (3a821 Thiamine mononitrate); 300 mg Vitamin B2 (Riboflavin); 3,000 mg Vitamin B3 (3a315 Niacinamide); 700 mg Vitamin B5 (3a841 Calcium-d-pantothenate); 240 mg Vitamin B6 (3a831 Pyridoxine hydrochloride); 1,400 mcg Vitamin B12 (Cynacobalamin); 80 mg Vitamin B9 (3a316 Folic acid); 16,000 mg Vitamin B4 (3a890 Choline chloride); 10,000 mcg Vitamin H (3a880 Biotin)

#### Technological additives

Antioxidants: 200 mg Citric acid (E330); 66.7 mg B.H.T (E321); 66.7 mg Propyl gallate (E310).

Anti-caking agents: 100 mg Colloidal silica (E551b)

#### Zootechnical additives

Digestibility enhancers: 40,000 FYT 6-Phytase (4a18); 22,400 TXU Endo-1,4- $\beta$ -xylanase (4a1602i); 10,000 TGU Endo-1,3(4)- $\beta$ -glucanase (4a1602i)

#### Coccidiostats

2,400 mg Salinomycin sodium (51776)

### **Description:**

Broilers have different feed requirements in terms of energy, proteins and minerals during different stages of their growth. It is important to adapt feed rations to these requirements for maximum production. Young broilers have a high protein requirement for the development of muscles, feathers, etc. As the broilers grow their energy requirements for the deposit of fat increase and their protein requirements decrease. DufaMix Broiler Starter is a premixture formulated to establish good appetite and achieve maximum early growth when included in corn and soybean meal based diets. An anticoccidial and enzymes are included for improved feed conversion, digestion and the reduction of wet droppings.

### **Inclusion rate:**

To be included in feed for approximately 5 to 12-day old broilers. The recommended inclusion rate for DufaMix Broiler Starter is 25 kg per tonne of feed.

**Storage conditions:**

Store dry, dark and below 25°C. Do not freeze.

**Packing:**

10 and 25 kg sacks.