

BROILER FINISHER

Corn-soybased diet

Analytical constituents:

74.3 % Crude ash	16.4 % Calcium	58.9 g/kg Lysine
0.19 % Crude fibre	0.2 % Magnesium	86.1 g/kg Methionine
14.4 % Crude protein	4.9 % Sodium	16.1 g/kg Threonine
0.02 % Crude fat	1.3 % Phosphorus	0.01 g/kg Tryptophan

Composition:

Calcium carbonate, Mono calcium phosphate, Sodium bicarbonate, Sodium chloride.

Additives (per kg)

Nutritional additives

Trace elements: 1,920 mg Fe (3b103 Iron(II)sulphate monohydrate); 64 mg I (3b202 Calcium iodate anhydrous); 480 mg Cu (3b504 Cupric(II)sulphate-pentahydrate); 2,240 mg Mn (3b502 Manganous(II) oxide); 2,240 mg Zn (3b603 Zinc oxide); 10 mg Se (3b801 Sodium Selenite)
Vitamins: 384,000 IU Vitamin A (3a672a Retinyl acetate); 96,000 IU Vitamin D3 (3a671); 1,280 mg Vitamin E (3a700i dl-tocopherol); 80 mg Vitamin K3 (3a711 Menadione nicotinamide bisulphite); 64 mg Vitamin B1 (3a821 Thiamine mononitrate); 192 mg Vitamin B2 (3a825ii Riboflavin); 480 mg Vitamin B5 (3a841 Calcium-d-pantothenate); 1,600 mg Vitamin B3 (3a315 Niacinamide); 160 mg Vitamin B6 (3a831 Pyridoxine hydrochloride); 960 mcg Vitamin B12 (Cynacobalamin); 48 mg Vitamin B9 (3a316 Folic acid); 9,600 mg Vitamin B4 (3a890 Choline chloride); 4,800 mcg Vitamin H (3a880 Biotin)

Zootechnical additives

Digestibility enhancers:

40,000 FYT 6-Phytase (4a18); 22,400 TXU Endo-1,4- β -xylanase (4a7); 10,000 TGU Endo-1,3(4)- β -glucanase (4a7)

Technological additives

Antioxidants: 160 mg Citric acid (E330); 66.7 mg B.H.T (E321); 66.7 mg Propyl gallate (E310)
Anti-caking agent: 119.4 g Clinoptilolite of sedimentary origin (1g568)
Emulsifying and stabilizing agent: 23.8 mg Glycerol polyethyleneglycol ricinoleate (E484)

Description:

Changes in body composition can be rapid during the finishing period and excessive fat deposition and loss of breast meat yield need to be carefully considered. DufaMix Broiler Finisher is formulated to complement corn and soybean meal based diets for optimal performance. Enzymes are included for improved feed conversion, digestion and the reduction of wet droppings.

Inclusion rate:

To be included in feed for broilers of approximately 28 days and older; 25 kg per tonne of feed.

Storage conditions:

Store dry, dark and below 25°C. Do not freeze.

Packing: 10 and 25 kg sacks.