

BROILER GROWER

Corn-soybased diet

Analytical constituents:

69.4 % Crude ash	14.8 % Calcium	72.3 g/kg Lysine
0.43 % Crude fibre	0.20 % Magnesium	101.1 g/kg Methionine
17.8 % Crude protein	5.00 % Sodium	25.8 g/kg Threonine
0.31 % Crude fat	4.20 % Phosphorus	0.01 g/kg Tryptophan

Composition:

Calcium carbonate, Mono calcium phosphate, Sodium bicarbonate, Sodium chloride.

Additives (per kg)

Nutritional additives

Trace elements: 2,400 mg Fe (3b103 Iron(II)sulphate monohydrate); 80 mg I (3b202 Calcium iodate anhydrous); 600 mg Cu (3b504 Cupric(II)sulphate-pentahydrate); 2,800 mg Mn (3b502 Manganous(II) oxide); 2,800 mg Zn (3b603 Zinc oxide); 10 mg Se (3b801 Sodium Selenite)
Vitamins: 400,000 IU Vitamin A (3a672a Retinyl acetate); 120,000 IU Vitamin D3 (3a671); 1,600 mg Vitamin E (3a700i dl-tocopherol); 100 mg Vitamin K3 (3a711 Menadione nicotinamide bisulphite); 80 mg Vitamin B1 (3a821 Thiamine mononitrate); 240 mg Vitamin B2 (3a825ii Riboflavin); 600 mg Vitamin B5 (3a841 Calcium-d-pantothenate); 2,000 mg Vitamin B3 (3a315 Niacinamide); 200 mg Vitamin B6 (3a831 Pyridoxine hydrochloride); 1,200 mcg Vitamin B12 (Cynacobalamin); 60 mg Vitamin B9 (3a316 Folic acid); 12,000 mg Vitamin B4 (3a890 Choline chloride); 6,000 mcg Vitamin H (3a880 Biotin)

Zootechnical additives

Digestibility enhancers:

40,000 FYT 6-Phytase (4a18); 22,400 TXU Endo-1,4- β -xylanase (4a7); 10,000 TGU Endo-1,3(4)- β -glucanase (4a7)

Coccidiostats

2,400 mg Salinomycin sodium (51766)

Technological additives

Antioxidants: 200 mg Citric acid (E330); 83.3 mg B.H.T (E321); 83.3 mg Propyl gallate (E310)
Anti-caking agent: 60 mg Colloidal silica (E551b); 34 g Clinoptilolite of sedimentary origin (1g568)
Emulsifying and stabilizing agent: 29.7 mg Glyceryl polyethyleneglycol ricinoleate (E484)

Description:

DufaMix Broiler Grower is formulated to complement corn and soybean meal based diets for optimal performance and to allow the bird to express its full meat yield potential. An anticoccidial and enzymes are included for improved feed conversion, digestion and the reduction of wet droppings.

Inclusion rate: included in feed for approximately 12 to 28-day old broilers; 25 kg per ton of feed.

Storage conditions: Store dry, dark and below 25°C. Do not freeze.

Packing: 10 and 25 kg sacks.